

**CLEAN VERSION OF CLAIMS**

**NUCLEIC ACID ANALYSIS USING COMPLETE N-MER ARRAYS**

Applicant: Kevin L. Gunderson et al.

Serial No.: 09/394,230

1. (Four Times Amended) A method of determining the presence of a mutation in a target polynucleotide, comprising the steps of:
- (a) providing at least two identical polynucleotide probe arrays, wherein each probe comprises a double stranded region and a single-stranded n-mer overhang region such that the overhangs in each array constitute a complete set of n-mers;
  - (b) hybridizing the target polynucleotide to said overhangs of probe polynucleotides in one array to generate a target hybridization pattern;
  - (c) hybridizing a reference polynucleotide to said overhangs of probe polynucleotides in a second array to generate a reference hybridization pattern; and
  - (d) determining the presence of a mutation in the target polynucleotide by comparing the reference and target hybridization patterns;
- wherein the method has a false positive rate of less than 1 per 3900 bp.
2. The method of claim 1, wherein in step b), the hybridized target polynucleotide is ligated to the probe.
3. The method of claim 1, wherein in step c), the hybridized reference polynucleotide is ligated to the probe.
4. The method of claim 1, wherein the overhangs have free 5'-ends.
5. The method of claim 1, wherein the overhangs have free 3'-ends.
6. The method of claim 1, wherein the n-mer comprises from about 4 to about 50 nucleotides.

7. The method of claim 1, wherein the mutation is a substitution mutation.
8. The method of claim 1, wherein the mutation is a deletion mutation.
9. (Once Amended) The method of claim 1, wherein the mutation is an insertion mutation.
10. The method of claim 1, in which said target polynucleotide is selected from the group consisting of: a cystic fibrosis transmembrane conductance regulator gene, a p53 gene, a mitochondrial DNA, or an HIV gene.

- 
- 03
11. (Once Amended) The method of claim 1, wherein the arrays are arranged in parallel.
  12. (Four Times Amended) A method of determining whether two or more target polynucleotides are identical, comprising the steps of:
    - (a) providing at least two identical polynucleotide probe arrays, wherein each probe comprises a double stranded region and a single-stranded n-mer overhang region such that the overhangs in each array constitute a complete set of n-mers, wherein each n-mer is at least 8 nucleotides in length;
    - (b) hybridizing first target polynucleotide to said overhangs of probe polynucleotides in one array to generate a first hybridization pattern;
    - (c) hybridizing second target polynucleotide to said overhangs of probe polynucleotides in a second array to generate a second hybridization pattern; and
    - (d) comparing the first and second hybridization patterns;wherein the method has a false positive rate of less than 1 per 3900 bp.

- 
13. The method of claim 12, wherein in step b), the hybridized target polynucleotide is ligated to the probe.
  14. The method of claim 12, wherein in step c), the hybridized reference polynucleotide is

AMENDMENT & RESPONSE UNDER 37 C.F.R. § 1.116 - EXPEDITED PROCEDURE

Serial Number: 09/394,230

Filing Date: September 13, 1999

Title: NUCLEIC ACID ANALYSIS USING COMPLETE N-MER ARRAYS

---

Page 12

Dkt: 1451.003US1

ligated to the probe.

15. The method of claim 12, wherein the overhangs have free 5'-ends.
  16. The method of claim 12, wherein the overhangs have free 3'-ends.
  17. The method of claim 12, wherein the n-mer comprises from about 4 to about 50 nucleotides.
  18. (Twice Amended) The method of claim 12, wherein the arrays are arranged in parallel.
-